

Technical Data Sheet

InVivoMAb anti-mouse CD4



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	BE0003-1
Clone:	GK1.5
Isotype:	Rat IgG2b, κ
Recommended Isotype Control(s):	InVivoMAb rat IgG2b isotype control, anti-keyhole limpet hemocyanin
Recommended Dilution Buffer:	InVivoPure pH 6.5 Dilution Buffer
Immunogen:	Mouse CTL clone V4
Reported Applications:	<i>in vivo</i> CD4 ⁺ T cell depletion Flow cytometry Western blot
Formulation:	PBS, pH 6.5 Contains no stabilizers or preservatives
Endotoxin:	$\leq 1\text{EU/mg}$ ($\leq 0.001\text{EU}/\mu\text{g}$) Determined by LAL gel clotting assay
Purity:	$\geq 95\%$ Determined by SDS-PAGE
Sterility:	0.2 μm filtration
Production:	Purified from cell culture supernatant in an animal-free facility
Purification:	Protein G
RRID:	AB_1107636
Molecular Weight:	150 kDa

Description

The GK1.5 monoclonal antibody reacts with the mouse CD4. The CD4 antigen is a 55 kDa cell surface type I membrane glycoprotein belonging to the immunoglobulin superfamily. CD4 acts as a co-receptor which in cooperation with the T cell receptor (TCR) interacts with class II MHC molecules displayed by antigen presenting cells (APC). CD4 is expressed by the majority of thymocytes, most helper T cells, a subset of NK-T cells and weakly by dendritic cells and macrophages. CD4 plays an important role in the development of T cells and is required for mature T cells to function optimally. The GK1.5 antibody has been shown to compete with clones YTS 177 and YTS 191 for CD4 binding.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration

experiment.


Application References

For a complete list of references, visit https://bioxcell.com/be0003-1?bxcs=9k1b3a#tab_references or scan the QR code below.



Binding Validation

Validation data shown below confirms that this clone binds to its target antigen. For lot specific binding validation data, e-mail technicalservice@bioxcell.com.



Bio X Cell, LLC
<https://bioxcell.com>
+1-866-787-3444
customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.
Not for resale.
Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC