Technical Data Sheet

InVivoMAb anti-human LFA-1α (CD11a)

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: **Lot Specific***
Volume: **Lot Specific***
Concentration: **Lot Specific* (generally 4 to 11 mg/ml) ***
Total Protein: **Lot Specific***

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	**BE0005**
Clone:	**TS-1/22.1.1.13**
Isotype:	Mouse IgG1
Recommended Isotype Control(s):	InVivoMAb mouse IgG1 isotype control, unknown specificity
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Immunogen:	Human cytolytic T cells
Reported Applications:	*in vitro* LFA-1 neutralization
Formulation:	PBS, pH 7.0 Contains no stabilizers or preservatives
Endotoxin:	≤1EU/mg (≤0.001EU/µg) Determined by LAL gel clotting assay
Purity:	≥95% Determined by SDS-PAGE
Sterility:	0.2 µm filtered
Production:	Purified from cell culture supernatant in an animal-free facility
Purification:	Protein G
RRID:	AB_1107580
Molecular Weight:	150 kDa

Description

The TS-1/22.1.1.13 monoclonal antibody reacts with human LFA-1α(lymphocyte function-associated antigen 1 alpha) also known as integrin alpha L chain and CD11a. LFA-1αand CD18 combine to form LFA-1 a 180 kDa glycoprotein and a member of the integrin family. LFA-1 is expressed on the surface of all leukocytes including lymphocytes monocytes macrophages and granulocytes. LFA-1 plays a central role in leukocyte intercellular adhesion through interactions with its ligands ICAM-1 (CD54) ICAM-2 (CD102) and ICAM-3 (CD50) and also functions in lymphocyte costimulatory signaling.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**
It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0005?bxcs=9k1b3a#tab_references or scan the QR code below.

Bio X Cell, LLC
https://bioxcell.com
+1-866-787-3444
customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.
Not for resale.
Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC