Technical Data Sheet

InVivoMAb anti-mouse CD22

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: **Lot Specific***
Volume: **Lot Specific***
Concentration: **Lot Specific* (generally 4 to 11 mg/ml) ***
Total Protein: **Lot Specific***

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	**BE0011**
Clone:	**Cy34.1**
Isotype:	Mouse IgG1, κ
Recommended Isotype Control(s):	InVivoMAb mouse IgG1 isotype control, unknown specificity
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Immunogen:	B10.D2 mouse splenocytes
Reported Applications:	*in vivo* B cell depletion in combination with anti-CD19 (clone 1D3) and anti-rat κ Light Chain (clone MAR 18.5) Flow cytometry Immunoprecipitation
Formulation:	PBS, pH 7.0 Contains no stabilizers or preservatives
Endotoxin:	≤1EU/mg (≤0.001EU/µg) Determined by LAL gel clotting assay
Purity:	≥95% Determined by SDS-PAGE
Sterility:	0.2 µm filtered
Production:	Purified from cell culture supernatant in an animal-free facility
Purification:	Protein G
RRID:	AB_1107613
Molecular Weight:	150 kDa

Description

The Cy34.1 monoclonal antibody reacts with mouse CD22, a member of the SIGLEC family of lectins. CD22 is expressed at high levels on the surface of mature follicular and marginal zone B lymphocytes, B-1 cells, and plasma cells and associates with the B-cell antigen receptor. CD22 mediates B cell adhesion to ligands on endothelial cells in the bone marrow. Additionally, CD22 negatively regulates B cell activation and prevents the development of autoimmune diseases. The Cy34.1 antibody has been shown to augment B cell proliferation in response to LPS or anti-mouse Ig µ chain.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**
It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration

experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0011?bxcs=9k1b3a#tab_references or scan the QR code below.

Bio X Cell, LLC
https://bioxcell.com
+1-866-787-3444
customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.
Not for resale.
Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC