# Technical Data Sheet

## InVivoMAb anti-mouse OX40 (CD134)

**Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.**

## Lot Specific Information

**Lot Number:** **Lot Specific***
**Volume:** **Lot Specific***
**Concentration:** **Lot Specific* (generally 4 to 11 mg/ml) ***
**Total Protein:** **Lot Specific***

*This information will be noted on the certificate of analysis that ships with this product.

## Product Information

| | |
|---|---|
| **Catalog Number:** | **BE0031** |
| **Clone:** | **OX-86** |
| **Isotype:** | Rat IgG1, κ |
| **Recommended Isotype Control(s):** | InVivoMAb rat IgG1 isotype control, anti-horseradish peroxidase |
| **Recommended Dilution Buffer:** | InVivoPure pH 7.0 Dilution Buffer |
| **Immunogen:** | Recombinant mouse OX40-CD4 chimeric protein |
| **Reported Applications:** | *in vivo* OX40 activation<br>*in vitro* OX40 activation<br>Western blot |
| **Formulation:** | PBS, pH 7.0<br>Contains no stabilizers or preservatives |
| **Endotoxin:** | ≤1EU/mg (≤0.001EU/μg)<br>Determined by LAL gel clotting assay |
| **Purity:** | ≥95%<br>Determined by SDS-PAGE |
| **Sterility:** | 0.2 μm filtration |
| **Production:** | Purified from cell culture supernatant in an animal-free facility |
| **Purification:** | Protein G |
| **RRID:** | AB_1107592 |
| **Molecular Weight:** | 150 kDa |

## Description

The OX-86 monoclonal antibody reacts with mouse OX-40 also known as CD134. OX-40 is a 50 kDa type I membrane glycoprotein and a member of the TNF receptor superfamily. OX-40 is expressed on activated CD4 and CD8 T cells, but is not found on resting naïve T cells or most resting memory T cells. Although it was originally thought that OX-40 expression was restricted to activated conventional T cells, it has now been visualized on activated regulatory T cells, NKT cells, NK cells, and neutrophils. OX-40 plays a major role in regulating both CD4 and CD8 T cell clonal expansion. It provides a costimulatory signal to an antigen-reacting naive T cells to prolong proliferation, as well as augment the production of several cytokines. This is demonstrated by OX-40 knockout mice which generate fewer primary effector CD4 T cells after immunization. Furthermore, in vivo treatment with an agonist antibody to OX-40 has been shown to strongly enhance the generation of antigen-specific effector T cells and prevent the induction of T cell tolerance. The OX-86 antibody is an agonistic antibody that has been shown to delay tumor growth in vivo.

## Storage

Store at the stock concentration at 4°C . **Do not freeze.**
It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

## Protocol Information


Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

## Application References

For a complete list of references, visit https://bioxcell.com/be0031?bxcs=9k1b3a#tab_references or scan the QR code below.

## Binding Validation

Validation data shown below confirms that this clone binds to its target antigen. For lot specific binding validation data, e-mail technicalservice@bioxcell.com.




**Bio X Cell, LLC**
https://bioxcell.com
+1-866-787-3444
customerservice@bioxcell.com

*Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.*
*Not for resale.*
**Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC**