Technical Data Sheet

InVivoMAb anti-mouse IL-10R (CD210)

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: **Lot Specific***
Volume: **Lot Specific***
Concentration: **Lot Specific* (generally 4 to 11 mg/ml) ***
Total Protein: **Lot Specific***

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	**BE0050**
Clone:	**1B1.3A**
Isotype:	Rat IgG1, κ
Recommended Isotype Control(s):	InVivoMAb rat IgG1 isotype control, anti-horseradish peroxidase
Recommended Dilution Buffer:	InVivoPure pH 6.5T Dilution Buffer
Immunogen:	Recombinant ligand-binding domain of mouse IL-10R
Reported Applications:	*in vivo* blocking of IL-10/IL-10R signaling *in vitro* blocking of IL-10R signaling Flow cytometry Western Blot
Formulation:	PBS, pH 6.5 0.01% Tween Contains no stabilizers or preservatives
Endotoxin:	≤1EU/mg (≤0.001EU/μg) Determined by LAL gel clotting assay
Purity:	≥95% Determined by SDS-PAGE
Sterility:	0.2 μm filtration
Production:	Purified from cell culture supernatant in an animal-free facility
Purification:	Protein G
RRID:	AB_1107611
Molecular Weight:	150 kDa

Description

The 1B1.3A monoclonal antibody reacts with mouse IL-10R (IL-10 receptor) also known as CD210. The IL-10R is a class II cytokine receptor and is expressed by a variety of cell types including thymocytes, T lymphocytes, B lymphocytes, NK cells, monocytes, and macrophages. Upon binding IL-10, IL-10R stimulation results in many pleiotropic, effects in immunoregulation and inflammation. IL-10R downregulates the expression of pro-inflammatory cytokines, MHC class II antigens, and co-stimulatory molecules on macrophages. It also enhances B lymphocyte survival, proliferation, and antibody production. IL-10R signaling can block NF-κB activity, and is involved in the regulation of the JAK-STAT signaling pathway. The 1B1.3A antibody is a neutralizing antibody and has been shown to block the binding of human IL-10, which cross-reacts with the mouse IL-10R. However, this clone does not recognize the human IL-10R. Bio X Cell is pleased to offer two recombinant, murine chimeric versions of the original 1B1.3A antibody, 1B1.3A-CP060 and 1B1.3A-CP073. The variable region sequences are identical to the original 1B1.3A but the constant region sequences have been switched from Rat IgG1, κ to mouse IgG2a, κ for use in murine models. Species-matched chimeric antibodies exhibit regulated effector functions—including Fc receptor binding and complement activation—and cause less immunogenicity and formation of anti-drug antibodies (ADAs) than xenogenic antibodies in animal models. Additionally, 1B1.3A-CP073 contains LALA-PG Fc-silencing mutations in the heavy chain rendering it unable to bind endogenous murine FcγR or C1q to induce antibody-

dependent, cell-mediated cytotoxicity (ADCC) or complement-dependent cytotoxicity (CDC). Antibodies with active Fc regions can engage immune cells via FcγRs, leading to the depletion of antigen expressing cells through mechanisms like ADCC or complement activation. Fc-silenced antibodies do not trigger these pathways and can block signaling without killing or depleting target cells. The highly controlled sequence and lack of genetic drift in recombinant antibodies provide more reliable and reproducible results over hybridoma derived antibodies.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**
It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

Protocol Information


Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit
https://bioxcell.com/be0050?bxcs=9k1b3a#tab_references
or scan the QR code below.

Binding Validation

Validation data shown below confirms that this clone binds to its target antigen. For lot specific binding validation data, e-mail technicalservice@bioxcell.com.



Bio X Cell, LLC
https://bioxcell.com
+1-866-787-3444
customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.
Not for resale.
Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC