Technical Data Sheet

InVivoMAb anti-mouse 4-1BBL (CD137L)

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: **Lot Specific***
Volume: **Lot Specific***
Concentration: **Lot Specific* (generally 4 to 11 mg/ml) ***
Total Protein: **Lot Specific***

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	**BE0110**
Clone:	**TKS-1**
Isotype:	Rat IgG2a, κ
Recommended Isotype Control(s):	InVivoMAb rat IgG2a isotype control, anti-trinitrophenol
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Immunogen:	Mouse 4-1BBL transfected NRK cells
Reported Applications:	*in vivo* 4-1BBL blockade ELISA
Formulation:	PBS, pH 7.0 Contains no stabilizers or preservatives
Endotoxin:	≤1EU/mg (≤0.001EU/µg) Determined by LAL gel clotting assay
Purity:	≥95% Determined by SDS-PAGE
Sterility:	0.2 µm filtered
Production:	Purified from cell culture supernatant in an animal-free facility
Purification:	Protein G
RRID:	AB_10949069
Molecular Weight:	150 kDa

Description

The TKS-1 monoclonal antibody reacts with mouse 4-1BB ligand (4-1BBL), a type II transmembrane glycoprotein also known as CD137L. 4-1BBL is a 97 kDa member of the TNF superfamily and is expressed by dendritic cells, macrophages, and activated B and T lymphocytes. Interaction of 4-1BBL with 4-1BB (CD137) provides costimulatory signals to both CD4 and CD8 T cells through the activation of NF-κB, c-Jun and p38 downstream pathways. The TKS-1 antibody has been shown to inhibit the binding of soluble 4-1BB to 4-1BBL in vitro.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**
It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0110?bxcs=9k1b3a#tab_references or scan the QR code below.

Bio X Cell, LLC
https://bioxcell.com
+1-866-787-3444
customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.
Not for resale.
Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC