Technical Data Sheet

InVivoMAb anti-human IL-4

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: **Lot Specific***
Volume: **Lot Specific***
Concentration: **Lot Specific* (generally 4 to 11 mg/ml) ***
Total Protein: **Lot Specific***

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	**BE0240**
Clone:	**MP4-25D2**
Isotype:	Rat IgG1, κ
Recommended Isotype Control(s):	InVivoMAb rat IgG1 isotype control, anti-horseradish peroxidase
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Immunogen:	Recombinant human IL-4 expressing CHO cells
Reported Applications:	*in vitro* IL-4 neutralization Flow cytometry
Formulation:	PBS, pH 7.0 Contains no stabilizers or preservatives
Endotoxin:	≤1EU/mg (≤0.001EU/µg) Determined by LAL gel clotting assay
Purity:	≥95% Determined by SDS-PAGE
Sterility:	0.2 µm filtered
Production:	Purified from cell culture supernatant in an animal-free facility
Purification:	Protein G
RRID:	AB_2687722
Molecular Weight:	150 kDa

Description

The MP4-25D2 monoclonal antibody reacts with human IL-4 (interleukin-4) a multifunctional 14 kDa cytokine. IL-4 is expressed primarily by activated Th2 cells and NK cells, and at lower levels by mast cells, and basophils. IL-4 signals through the IL-4Rα. Upon receptor binding IL-4 stimulates activated B and T lymphocyte proliferation, and the differentiation of B cells into plasma cells. It also induces B cell class switching to IgE, and up-regulates MHC class II production while decreasing the production of Th1 cells, macrophages, IFNγ, and dendritic cell IL-12. Like other Th2 associated cytokines, IL-4 is involved in the airway inflammation observed in the lungs of patients with allergic asthma. The MP4-25D2 monoclonal antibody has been shown to neutralize the bioactivity of natural or recombinant human IL-4.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**
It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration

experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0240?bxcs=9k1b3a#tab_references or scan the QR code below.

Bio X Cell, LLC
https://bioxcell.com
+1-866-787-3444
customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.
Not for resale.
Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC