Technical Data Sheet

InVivoMAb recombinant Flt-3L-Ig (hum/hum)

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: **Lot Specific***
Volume: **Lot Specific***
Concentration: **Lot Specific* (generally 4 to 11 mg/ml) ***
Total Protein: **Lot Specific***

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	**BE0342**
Clone:	**Flt-3L Fc-G1**
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Formulation:	PBS, pH 7.0 Contains no stabilizers or preservatives
Endotoxin:	≤1EU/mg (≤0.001EU/µg) Determined by LAL gel clotting assay
Purity:	≥95% Determined by SDS-PAGE
Sterility:	0.2 µm filtration
Production:	Purified from cell culture supernatant in an animal-free facility
Purification:	Protein A
RRID:	BE0342

Description

Flt-3L (FMS-related Tyrosine Kinase 3 Ligand) is an endogenous protein that functions as a cytokine and growth factor. Flt-3L is crucial for the development of conventional dendritic cells (cDCs) and plasmacytoid dendritic cells (pDCs). Recombinant Flt-3L-Ig is a fusion protein consisting of human Flt-3L fused to the Fc portion of human IgG1. This fusion protein is useful for activating Flt3 signaling and inducing the expansion of DC populations. Human Flt-3L-Ig is frequently reported to stimulate Flt3 signaling in vivo in mice.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**
It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0342?bxcs=9k1b3a#tab_references or scan the QR code below.

Bio X Cell, LLC
https://bioxcell.com
+1-866-787-3444
customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.
Not for resale.
Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC