Technical Data Sheet

InVivoPlus anti-mouse GITR

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: **Lot Specific***
Volume: **Lot Specific***
Concentration: **Lot Specific* (generally 4 to 11 mg/ml) ***
Total Protein: **Lot Specific***

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	**BP0063**
Clone:	**DTA-1**
Isotype:	Rat IgG2b, λ
Recommended Isotype Control(s):	InVivoPlus rat IgG2b isotype control, anti-keyhole limpet hemocyanin
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Immunogen:	Mouse CD25+ CD4+ T cells
Reported Applications:	*in vivo* GITR stimulation
Formulation:	PBS, pH 7.0 Contains no stabilizers or preservatives
Endotoxin:	≤0.5EU/mg (≤0.0005EU/μg) Determined by LAL gel clotting assay
Purity:	≥95% Determined by SDS-PAGE
Sterility:	0.2 μm filtration
Production:	Purified from cell culture supernatant in an animal-free facility
Purification:	Protein G
Aggregation:	<5% Determined by SEC
RRID:	AB_1107688
Molecular Weight:	150 kDa

Murine Pathogen Test Results

Mouse Norovirus: Negative, Mouse Parvovirus: Negative, Mouse Minute Virus: Negative, Mouse Hepatitis Virus: Negative, Reovirus Screen: Negative, Lymphocytic Choriomeningitis virus: Negative, Lactate Dehydrogenase-Elevating Virus: Negative, Mouse Rotavirus: Negative, Theiler's Murine Encephalomyelitis: Negative, Ectromelia/Mousepox Virus: Negative, Hantavirus: Negative, Polyoma Virus: Negative, Mouse Adenovirus: Negative, Sendai Virus: Negative, Mycoplasma Pulmonis: Negative, Pneumonia Virus of Mice: Negative, Mouse Cytomegalovirus: Negative, K Virus: Negative

Description

The DTA-1 monoclonal antibody reacts with mouse GITR (glucocorticoid-induced TNFR-related gene), a 66-70 kDa co-stimulatory immune checkpoint molecule belonging to the Tumor Necrosis Factor superfamily (TNFRSF18). GITR is expressed at low levels on resting T lymphocytes and at high levels on regulatory T cells. GITR is upregulated on activated T cells where it provides co-stimulation. GITR ligand (GITRL) is found on B cells, macrophages, dendritic and endothelial cells, and is implicated in regulating both innate and adaptive immune responses. GITR is also thought to play a key role in dominant immunological self-tolerance maintained by regulatory T cells. Knockout studies in mice also suggest the role of this receptor is in the regulation of CD3-driven T cell activation and programmed cell death. The DTA-1 antibody is an agonistic antibody that is commonly used to induce GITR signaling in vivo.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**
It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/bp0063?bxcs=9k1b3a#tab_references or scan the QR code below.

Bio X Cell, LLC
https://bioxcell.com
+1-866-787-3444
customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.
Not for resale.
Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC